

The following is an excerpt from Chapter 10 of *Calculus, Early Transcendentals* by James Stewart. It has been reformatted slightly from its appearance in the textbook so that it corresponds to the LaTeX article style.

## 1 The Binomial Series

You may be acquainted with the Binomial Theorem, which states that if  $a$  and  $b$  are any real numbers and  $k$  is a positive integer, then

$$\begin{aligned} (a + b)^k &= a^k + ka^{k-1}b + \frac{k(k-1)}{2!}a^{k-2}b^2 + \frac{k(k-1)(k-2)}{3!}a^{k-3}b^3 \\ &+ \cdots + \frac{k(k-1)(k-2)\cdots(k-n+1)}{n!}a^{k-n}b^n \\ &+ \cdots + kab^{k-1} + b^k \end{aligned}$$

The traditional notation for the binomial coefficient is

$$\binom{k}{0} = 1 \quad \binom{k}{n} = \frac{k(k-1)(k-2)\cdots(k-n+1)}{n!} \quad n = 1, 2, \dots, k$$

which enables us to write the Binomial Theorem in the abbreviated form

$$(a + b)^k = \sum_{n=0}^k \binom{k}{n} a^{k-n} b^n$$

In particular, if we put  $a = 1$  and  $b = x$ , we get

$$(1 + x)^k = \sum_{n=0}^k \binom{k}{n} x^n \tag{1}$$

One of Newton's accomplishments was to extend the Binomial Theorem (Equation 1) to the case where  $k$  is no longer a positive integer. In this case the expression for  $(1 + x)^k$  is no longer a finite sum; it becomes an infinite series.

Assuming that  $(1 + x)^k$  can be expanded as a power series, we compute its Maclaurin series in the usual way:

$$\begin{array}{ll} f(x) = (1 + x)^k & f(0) = 1 \\ f'(x) = k(1 + x)^{k-1} & f'(0) = k \\ f''(x) = k(k-1)(1 + x)^{k-2} & f''(0) = k(k-1) \\ f'''(x) = k(k-1)(k-2)(1 + x)^{k-3} & f'''(0) = k(k-1)(k-2) \\ \vdots & \vdots \\ f^{(n)}(x) = k(k-1)\cdots(k-n+1)(1 + x)^{k-n} & f^{(n)}(0) = k(k-1)\cdots(k-n+1) \end{array}$$

$$(1+x)^k = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = \sum_{n=0}^{\infty} \frac{k(k-1)\cdots(k-n+1)}{n!} x^n$$

If  $u_n$  is the  $n$ th term of this series, then

$$\begin{aligned} \left| \frac{u_{n+1}}{u_n} \right| &= \left| \frac{k(k-1)\cdots(k-n+1)(k-n)x^{n+1}}{(n+1)!} \cdot \frac{n!}{k(k-1)\cdots(k-n+1)x^n} \right| \\ &= \frac{|k-n|}{n+1} |x| = \frac{1 - \frac{k}{n}}{1 + \frac{1}{n}} |x| \rightarrow |x| \quad \text{as } n \rightarrow \infty \end{aligned}$$

Thus, by the Ratio Test, the binomial series converges if  $|x| < 1$  and diverges if  $|x| > 1$ .

**Definition 1 (The Binomial Series).** If  $k$  is any real number and  $|x| < 1$ , then

$$\begin{aligned} (1+x)^k &= 1 + kx + \frac{k(k-1)}{2!}x^2 + \frac{k(k-1)(k-2)}{3!}x^3 + \cdots \\ &= \sum_{n=0}^{\infty} \binom{k}{n} x^n \end{aligned}$$

where

$$\binom{k}{n} = \frac{k(k-1)\cdots(k-n+1)}{n!} \quad (n \geq 1) \quad \binom{k}{0} = 1$$

We have proved Definition 1 under the assumption that  $(1+x)^k$  has a power series expansion. For a proof without that assumption see Exercise 21.

Although the binomial series always converges when  $|x| < 1$ , the question of whether or not it converges at the endpoints,  $\pm 1$ , depends on the value of  $k$ . It turns out that the series converges at 1 if  $-1 < k \leq 0$  and at both endpoints if  $k \geq 0$ . Notice that if  $k$  is a positive integer and  $n > k$ , then the expression for  $\binom{k}{n}$  contains a factor  $(k-k)$ , so  $\binom{k}{n} = 0$  for  $n > k$ . This means that the series terminates and reduces to the ordinary Binomial Theorem (Equation 1) when  $k$  is a positive integer.

Although, as we have seen, the binomial series is just a special case of the Maclaurin series, it occurs frequently and so it is worth remembering.

**Example 1.** Expand  $\frac{1}{(1+x)^2}$  as a power series.

*Solution.* We use the binomial series with  $k = -2$ . The binomial coefficient is

$$\begin{aligned} \binom{-2}{n} &= \frac{(-2)(-3)(-4)\cdots(-2-n+1)}{n!} \\ &= \frac{(-1)^n 2 \cdot 3 \cdot 4 \cdots n(n+1)}{n!} = (-1)^n (n+1) \end{aligned}$$

and so, when  $|x| < 1$ ,

$$\begin{aligned} \frac{1}{(1+x)^2} &= (1+x)^{-2} = \sum_{n=0}^{\infty} \binom{-2}{n} x^n \\ &= \sum_{n=0}^{\infty} (-1)^n (n+1) x^n \end{aligned} \quad \square$$

**Example 2.** Find the Maclaurin series for the function  $f(x) = 1/\sqrt{4-x}$  and its radius of convergence.

*Solution.* As given,  $f(x)$  is not quite of the form  $(1+x)^k$  so we rewrite it as follows:

$$\frac{1}{\sqrt{4-x}} = \frac{1}{\sqrt{4(1-\frac{x}{4})}} = \frac{1}{2\sqrt{1-\frac{x}{4}}} = \frac{1}{2} \left(1 - \frac{x}{4}\right)^{-1/2}$$

Using the binomial series with  $k = -\frac{1}{2}$  and with  $x$  replaced by  $-x/4$ , we have

$$\begin{aligned} \frac{1}{\sqrt{4-x}} &= \frac{1}{2} \left(1 - \frac{x}{4}\right)^{-1/2} = \frac{1}{2} \sum_{n=0}^{\infty} \binom{-\frac{1}{2}}{n} \left(-\frac{x}{4}\right)^n \\ &= \frac{1}{2} \left[ 1 + \binom{-\frac{1}{2}}{1} \left(-\frac{x}{4}\right) + \frac{\binom{-\frac{1}{2}}{2} \left(-\frac{3}{2}\right)}{2!} \left(-\frac{x}{4}\right)^2 + \frac{\binom{-\frac{1}{2}}{3} \left(-\frac{3}{2}\right) \left(-\frac{5}{2}\right)}{3!} \left(-\frac{x}{4}\right)^3 \right. \\ &\quad \left. + \dots + \frac{\binom{-\frac{1}{2}}{n} \left(-\frac{3}{2}\right) \left(-\frac{5}{2}\right) \dots \left(-\frac{1}{2} - n + 1\right)}{n!} \left(-\frac{x}{4}\right)^n + \dots \right] \\ &= \frac{1}{2} \left( 1 + \frac{1}{8}x + \frac{1 \cdot 3}{2! 8^2}x^2 + \frac{1 \cdot 3 \cdot 5}{3! 8^3}x^3 + \dots + \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-1)}{n! 8^n}x^n + \dots \right) \end{aligned}$$

We know from Definition 1 that this series converges when  $|-x/4| < 1$ , that is,  $|x| < 4$ , so the radius of convergence is  $R = 4$ .  $\square$